

THE SUITABILITY OF EXPANDED SHALE AS AN AMENDMENT FOR CLAY SOILS

John J. Sloan¹, Steve W. George², Wayne A. Mackay³, Phil Colbaugh³, and Sam Feagley²

Texas A&M University Research and Extension Center – Dallas

17360 Coit Road

Dallas, TX 75252

¹ Assistant Professor, Texas Agricultural Experiment Station

² Professor, Texas Cooperative Extension

³ Associate Professor, Texas Agricultural Experiment Station

THE SUITABILITY OF EXPANDED SHALE AS AN AMENDMENT FOR CLAY SOILS

SUMMARY

Excessive soil moisture in clay soils can cause poor aeration and adversely affect plant growth. Small and large diameter expanded shales (ExSh), quartz sand, sphagnum peat moss (SPM), and cottonseed hulls (CH) were evaluated as soil amendments for Austin silty clay soil. A 3 inch (7.5 cm) layer of each amendment was incorporated to a depth of 6 inches (15 cm), resulting in a 1:1 mixture by volume. Pansies (*Viola x wittrockiana* 'Crown Azure Blue') were grown from December to June, followed by scaevola (*Scaevola aemula* 'New Wonder') from June to November for two growing seasons. Foliage quality and extent of flowering were evaluated bi-weekly. Pansy root weights and above-ground biomass were quantified at the end of each growing season. None of the amendments significantly affected pansy foliage quality or the number of blooms per plant. Small diameter ExSh and SPM decreased pansy N content the first year after application, but not the second. During the first growing season, when soils were frequently saturated due to excessive rainfall, pansy root weights were significantly higher in soils amended with the small and large diameter ExSh. Large diameter ExSh treatments significantly increased the survival rate of transplanted scaevola plants and also the quality of foliage and percent blossom coverage during both growing seasons. Cottonseed hulls also increased scaevola survival for both growing seasons, but did not consistently improve scaevola foliage quality or bloom coverage. Of the five amendments tested, large diameter ExSh consistently improved overall plant performance more than the other amendments.

Fine-textured clay soils, such as those in the Blackland region of Texas, present unique problems for maintaining healthy plant growth because they exhibit low permeability to air and water (Puentes et al, 1988). During periods of high soil moisture, such as following rainfall or irrigation, there is a reduction in air-filled pore volume (Grable and Siemer, 1968; Jayawardane and Meyer, 1985). Furthermore, the obstruction of pores by infiltrating water hinders air movement into the soil (Erickson, 1982). When water is added to soil during warm temperatures, oxygen is decreased even further due to increased microbial and root respiration (Hendrix et al., 1988; McAfee et al., 1989). The combined effect of these processes is that soil oxygen levels decrease following water infiltration into the soil (Dowdell et al., 1979; Meek et al., 1983).

Root elongation is directly affected by soil aeration (Voorhees et al., 1975). In the absence of oxygen in the rooting zone, cell elongation ceases within two to three minutes and root tips will start to die if oxygen is not restored within thirty minutes (Huck, 1970). If low oxygen conditions persist, such as in water-saturated clay soils, root growth will be stunted and the plant may be more susceptible to disease. Since roots are directly responsible for uptake of water and minerals, which are essential for plant growth and development (Salisbury and Ross, 1985), poor root development will decrease the quantity and quality of above-ground plant foliage. Soil amendments that increase porosity in fine textured soils might improve plant growth and quality.

Urban dwellers living on fine-textured clay soils could benefit from soil amendments that improve drainage in their garden and flower beds. Sphagnum peat moss (SPM) is an organic material composed of partially decomposed sphagnum moss (*Sphagnum spp.*). It has long been used as a component in container growing mediums and in US Golf Association putting greens (Carlson et al., 1998). In sand-based growing media, SPM improved turf quality more than several inorganic amendments (Bigelow et al., 2001), but its ability to improve the quality of flowering plants in clay soil is not documented. Many gardeners intuitively believe that sand will improve drainage in a clay soil. Quartz sand is used to construct golf putting greens (USGA, 1993) and has been investigated as a top dressing for sports fields (Baker and Canaway, 1992). However, little is known of its effect on flowering plants when mixed into a clay textured soil. Cottonseed hulls, a byproduct of the cotton ginning industry, contain relatively high

amounts of fiber and some protein and are usually used for cattle feed (Hsu et al., 1986). The coarse-textured characteristic of CH may increase drainage and aeration when mixed with a clay-textured soil. Also, decomposing CH in the soil may provide a source of nitrogen and micronutrients.

Expanded shale (ExSh) is a lightweight, porous, inorganic material commonly used as an ingredient in cement for construction and road surfacing. It is produced by heating mined shale to a temperature of 2200°F (1200°C). Individual ExSh aggregates are able to absorb 761 lb·ton⁻¹ water (380 kg·t⁻¹), of which over half is readily available to plants (Sloan et al., 2000). The heating process creates calcium and magnesium oxides on the surface of ExSh aggregates, which generate an alkaline pH (8-10) when measured in water (Sloan et al., 2000). However, ExSh has a low calcium carbonate equivalent (<0.5%) and will not measurably affect the pH of neutral or calcareous soils. Expanded shale has no significant cation exchange capacity, but the presence of oxides allows it to absorb significant amounts of soluble phosphorus (Sloan et al., 2000). Due to its relatively inert chemical nature, ExSh has little direct effect on soil chemical properties. However, it could have indirect effects on soil chemistry and biology by modifying the soil physical environment.

We hypothesized that ExSh would improve plant growth in clay soils, more so than sand or organic amendments, by increasing soil porosity, thereby increasing internal water drainage and oxygen supply to roots. The objective of this research was to determine the effect of ExSh amendments, sand, SPM and CH, on growth and performance of winter-grown pansies and summer-grown scaevola in a poorly-drained clay soil.

Materials and Methods

An Austin silty clay soil (fine-silty, carbonatic, thermic Udorthentic Haplustolls) at the Texas A&M University Research and Extension Center at Dallas was amended with two diameter sizes of ExSh, quartz sand, SPM, and CH. The experimental design was a randomized complete block comprised of six blocks. Each amendment was applied as a 3 inch (7.5 cm) layer on the surface of a 3x3 ft (91x91 cm) plot and then hand-incorporated with a spade into the upper 3 inches (7.5 cm) of soil, resulting in approximately 6 inches (15 cm) of 1:1 amended soil by volume. Six treatments were included in the

study. Unamended soil served as a control (Ctrl). For the control, 3 inches (7.5 in) of native soil from an adjacent area was added to each plot and incorporated in a similar fashion to the other amendments. The primary focus of the study was on a 0.039 to 0.118 inches (1 to 3 mm) small-diameter ExSh aggregate (SSh) and a 0.118 to 0.354 inch (3 to 6 mm) large-diameter ExSh aggregate (LSH) (Texas Industries, Dallas, Texas). Sphagnum peat moss (Sun Gro Horticulture, Inc, Bellevue, Washington) and coarse quartz sand (ACME Materials, Inc, Allen, Texas) represent two forms of soil amendments frequently added to urban soils to improve tilth and drainage. Cottonseed hulls (Martindale Feed Mill, Valley View, Texas) were also evaluated as a potential soil amendment.

For the next two years, *Viola x witrockiana* 'Crown Azure Blue' were grown in the amended soils from December to June, followed by fan flower, *Scaevola aemula* 'New Wonder', from June to November. *Scaevola* is a native Australian plant that typically performs poorly in clay-textured soils (Bhalla and Sweeney 1998; Fischer 1990, 1991; Gerlach 1990). During each respective season, each plot was planted with either four pansy plants or two *scaevola* plants. Pansy plants were evaluated biweekly by rating foliage quality on a scale of one to ten, with one representing worst and ten representing best, and by counting the number of open blossoms per plant. *Scaevola* foliage was evaluated in a similar fashion to the pansies, but instead of counting *scaevola* blossoms, they were ranked on a scale of zero to ten, with zero representing no blossoms and ten representing maximum blossom coverage. Prior to each planting, a complete fertilizer formulation (21N-3.1P-11.6K) (Calloway's Premium Lawn Fertilizer, Plano, TX) was added to each plot at a rate of 178, 27, and 98 lb·acre⁻¹ (200, 30, and 110 kg ha⁻¹) of N, P, and K, respectively. During the growing season, Peters All purpose Plant Food (20N-8.8P-16.6K; Grace-Sierra Horticultural Products Co., Milpitas, California) was applied once per month at N, P, and K rates of 24.5, 10.8, and 20.2 lb·acre⁻¹ (28, 12, and 23 kg·ha⁻¹), respectively. Sphagnum peat moss and CH also added nitrogen to the soil in amounts of 140 and 150 lb acre⁻¹ (160 and 180 kg ha⁻¹), respectively. However, with C:N ratios of 60 for SPM and 100 for CH, the nitrogen was not immediately available for plant uptake. Irrigation was applied by hand. The frequency and quantity of irrigation was based on visual observations of soil moisture in the top 3 inches (7.5 cm). When irrigation was applied, all plots received approximately the same amount of water.

Pansies were harvested at the end of each growing season to determine above-ground dry matter yield, total nitrogen content, and root weights. Above-ground plant tissue was dried at 140°F (60°C), weighed, and then ground to pass a 40 mesh sieve. Total N was measured using a micro-kjeldahl acid digestion followed by steam distillation (Bremner, 1996). At the end of each season, the root ball from each plant was extracted from the soil with a straight-tined pitchfork and washed with water to remove soil. During the first year, the original root ball was clearly visible, so only roots extending beyond the original root ball were removed for quantification. For the second year, it was difficult to identify the original root ball, so the entire root weight, including the original root ball, was quantified. Therefore, root weights were generally larger the second year. Roots were dried at 140°F (60°C) for 48 h before weighing.

Since our primary focus was on the effect of the inorganic ExSh amendments, we allowed the soil amendments to physically equilibrate with the soil for seven months before measuring soil bulk density. Soil cores of known volume [3-inch diameter x 2-inch depth (7.5-cm x 5-cm)] were extracted from the 2 to 4 inch (5 to 10 cm) depth of each plot using a hammer-driven cylindrical core sampler. The inner cylinder consisted of three consecutive 2-inch (5-cm) sections. Soil was trimmed flush with the top and bottom of the middle cylinder section. Cores were dried at 221°F (105°C) for 48 h and then weighed. Bulk density was calculated as the oven dry soil weight divided by the cylinder volume (Blake and Hartge, 1986).

Data was analyzed using the Statistical Analysis Software (SAS Institute, 2001). For qualitative pansy and scaevola ratings, a univariate analysis showed that the data had a normal distribution. Therefore, data was analyzed by analysis of variance (ANOVA) with seasonal treatment means separated by Duncan's Multiple Range Test (DMRT). Plant survival counts were transformed using an arcsine function before performing an ANOVA and separation of treatment means by DMRT. Soil bulk density, pansy yield, nitrogen content, and root weights were analyzed by an ANOVA with treatment means separated by DMRT.

Results and Discussion

Growing conditions: Fig. 1A shows monthly rainfall distribution and average monthly maximum temperatures for 1997 and 1998 when this study was conducted. In North Central Texas, the pansy growing season generally corresponds to the period from October to May and the scaevola growing season from May to November. Rainfall during the first half of 1997, which represents the bulk of the pansy growing season, was generally high. Rainfall during the same period of 1998 was more moderate. Rainfall during the latter half of each year, which corresponds to the scaevola growing season, tended to be sporadic. However, July and August of 1998 were particularly dry and research plots required frequent irrigation.

Average monthly maximum temperatures were similar during both years (Fig. 1B), but in 1998, temperatures from May to Aug were consistently higher than in 1997. This period corresponds to the end of the pansy growing season and the beginning of the scaevola growing season. In general, temperature was not a major factor during the two-year study. For pansy growth, the 1997 growing season was wetter than 1998. For scaevola growth, no single year was more favorable for growth than the other.

Soil Bulk Density: Bulk density is a good indicator of potential rooting problems due to overly-compacted soil. Jones (1983) found that for clay-textured soils, similar to the Austin silty clay from the present study, root activity of several crops was greatest at soil bulk densities in the range of 1100 to 1300 kg·m⁻³ (68.5 to 81.0 lb·ft³). Figure 2 shows the effect of soil amendments on bulk density of the Austin silty clay soil measured seven months after application. Bulk density of the unamended clay soil (Ctrl) was approximately 900 kg·m⁻³ (56.1 lb·ft³), which is typical for a recently tilled clay-textured soil (Logsdon et al., 1999; Logsdon and Cambardella, 2000). The small and large diameter ExSh treatments (SSh and LSh) had no significant effect on soil bulk density. Sphagnum peat moss and CH both significantly decreased soil bulk density. The quartz sand treatment was the only amendment that increased bulk density of the clay soil. None of the measured bulk densities were high enough to present a barrier to root growth.

Plant Survival: During the first growing season, pansy plant survival was at or near 100% for all treatments (Table 1). For the second growing season, plant survival for the control decreased to 71%, but there were no significant differences between the control and amended plots. In regards to pansy survival, ExSh treatments performed the same as other soil amendments.

The effect of ExSh amendments, in particular LSh, on scaevola survival was more dramatic, especially during the first year after application (Table 1). Large diameter ExSh and CH significantly increased plant survival relative to the control during both growing seasons. Small diameter ExSh and SPM also significantly increased scaevola survival during the second growing season.

Plant Quality: The primary function of ornamental plants is to be aesthetically pleasing. For flowering annuals, this is primarily determined by the appearance of foliage and flowers. None of the soil amendments tested in this study had a major influence on pansy foliage quality relative to the unamended control (Fig. 3A), except for CH, which significantly lowered pansy foliage quality during the 1996-97 growing season. During the 1996-97 growing season, SPM was the only amendment that had higher blossom ratings than the control (Fig. 3B). Large diameter ExSh had no effect on pansy foliage production and flowering, but SSh, sand, and CH actually decreased pansy foliage quality and number of blossoms relative to the control during the 1996-97 season. During the 1997-98 growing season, pansy foliage ratings and blossom numbers were nearly identical for the control and all soil amendments.

Soil amendments had an obvious beneficial effect on the quality of scaevola foliage and blossom coverage (Fig. 4). Large and small diameter ExSh and SPM increased the quality of scaevola foliage (Fig. 4A) and extent of flowering (Fig. 4B) relative to the control during both the 1996-97 and 1997-98 growing seasons. Cottonseed hulls significantly increased scaevola foliage quality in 1998 (Fig 4A) and scaevola blossom coverage in 1997 (Fig. 4B). Scaevola is a native Australian plant that grows better in well drained containers than in poorly-drained clay-textured soils (Steffen, 1989). It is likely that the ExSh and SPM amendments improved the scaevola rooting environment, which translated to improved above-ground growth. The improved rooting environment might be attributed to better soil aeration, more favorable soil moisture levels, or possibly, a combination of these factors.

Pansy yield and nitrogen content: Total above-ground biomass production can be an indication of how soil amendments affect plant growth. Biomass data showed that the effects of soil amendments on pansy growth were either neutral or negative during the first year after application, but by the second year, no negative effects were observed (Fig. 3C). During the 1996-97 growing season, only LSh and SPM had dry matter yields equivalent to the control. All other amendments reduced dry matter yield relative to the control. The trend among treatments for 1996-97 is nearly identical to that for foliage ratings and blossom numbers (Fig. 3A,B), which demonstrates that plant quality is closely allied to overall above-ground plant biomass. During the 1997-98 growing season, none of the soil amendments significantly affected dry matter yield compared to the control.

Cottonseed hulls was the only amendment that did not significantly decrease plant N content during either growing season (Fig. 3D). Large diameter ExSh and sand decreased plant N the second growing season, but not the first. Sphagnum peat moss significantly decreased pansy nitrogen content during both growing seasons, probably due to immobilization of soil N by soil microorganisms after the addition of a readily available carbon source. Small diameter ExSh also decreased pansy N content during both seasons, but not as much as SPM during the 1997-98 growing season. For the most part, the soil amendments had no major influence on pansy nitrogen content. The nitrogen content for all treatments was near the lower limit of the range typically required for optimal plant growth (Marschner, 1986).

Pansy Root Weights: Pansy root data must be interpreted in light of the fact that data for the first year includes only roots extending beyond the original root ball whereas data for the second year includes the entire root weight. Therefore, we focus only on differences among treatments within each year and do not compare root weights between years. At the end of the 1996-97 season, pansy root weights for the SSh and LSh treatments were significantly greater than the control and all other soil amendments (Fig. 3E). Climatic data showed that rainfall from February to May during the 1996-97 pansy growing season [15.6 inches (39.6 cm)] was considerably higher than the same period during the 1997-98 growing season [8.1 inches (20.5 cm)]. The Austin silty clay soil is poorly drained, so high amounts of precipitation can lead to periods of excessive soil moisture. High soil moisture reduces soil aeration,

which may contribute to poor root development and increased root disease. It is possible that in the 1996-97 growing season, excess soil water drained more easily from soils amended with ExSh than from other soil amendments and this created a more favorable environment for the production and proper functioning of root tissue. Pansy root weights for the 1997-98 growing season were not significantly affected by soil amendments. Excessive soil moisture was probably not a problem during the 1997-98 growing season due to less rainfall.

The ability of soil amendments to improve plant performance depends on the type of plant and on prevailing weather conditions. Plants that have a low tolerance for poor soil aeration, such as scaevola, may perform better in soils amended with LSh [0.118 to 0.354 inches (3 to 6 mm)] and SPM. Small diameter ExSh [0.039 to 0.118 inches (1 to 3 mm)] had no consistent effect on pansy or scaevola growth. During periods of excessive rainfall, plant root growth in clay soils may be aided by the incorporation of LSh. Cottonseed hulls tended to decrease plant performance the first year they were applied, probably due to nitrogen immobilization induced by a high C:N ratio, but improved it the second season. In terms of overall plant performance, the positive responses from LSh were more consistent than the other amendments evaluated. However, in the short term (1 to 2 years), a single application of SPM also produced positive responses in plant performance. Incorporation of LSh into soil is essentially a permanent modification of the soil's physical properties. It may be feasible to use this practice in highly visible and intensively utilized planting beds.

ACKNOWLEDGMENTS

This research was partially supported with funding from Texas Industries, Dallas, Texas. Valuable statistical guidance was provided by Dr. Bill Raun (Oklahoma State University) and Dr. Jim Heitholt (Texas Agricultural Experiment Station).

Literature Cited

- Baker, S.W. and Canaway, P.M. 1992. The effect of sand top dressing on the performance of winter games pitches of different construction types: play quality. *J. Sports Turf Res. Inst.* 68:62-72.
- Bhalla, P.L. and K. Sweeney. 1998. Micropropagation of scaevola – Australian native of ornamental horticulture. *Australian J. Exp. Agric.* 38:399-401,
- Bigelow, C.A., D.C. Bowman, D.K. Cassel, and T.W. Ruffy, Jr. 2001. Creeping bentgrass response to inorganic soil amendments and mechanically induced subsurface drainage and aeration. *Crop Sci.* 41:797-805.
- Blake, G.R. and K.H. Hartge. 1986. Bulk density. pp. 363-375. In: A. Klute (ed). *Methods of Soil Analysis: Part 1—Physical and Mineralogical Methods.* 2nd ed. Soil Sci. Soc. Am. Book Ser. No. 5. Am. Soc. Agron. and Soil Sci. Soc. Amer., Inc. Madison, Wisc.
- Bremner, J.M. 1996. Nitrogen—total. pp.1085-1121. In: D.L. Sparks et al. (ed.) *Methods of soil analysis. Part 3 Chemical methods.* Soil Sci. Soc. Amer. Book Ser. 5. Soil Science Society of America. Madison, Wis.
- Carlson, M.S. , C.L. Kerkman, and W.R. Kussow. 1998. Peats and supplements for root zone mixes. *Golf Course Management* 66(9):70-74.
- Dowdell, R.J., R. Crees, J.R. Burford, and R.Q. Cannell. 1979. Oxygen concentrations in a clay soil after ploughing or direct drilling. *J. Soil Sci.* 30:239-245.

- Erickson, A.E. 1982. Tillage effects on soil aeration. pp. 91-104. In: Predicting Tillage Effects on Soil Physical Properties and Processes. Amer. Soc. Agron. Special Pub. 44 American Society of Agronomy. Madison, Wis.
- Fischer, P. 1990. Effect of pH value and phosphate surplus. *Zierpflanzenbau* 30:76-78.
- Fischer, P. 1991. Wirkamseit von Eisenchelaten. *Deutscher Gartenbau*. 13:816-819.
- Gerlach, W.W.P. 1990. Verticillium-Welke als neue Krankheit an Scaevola. *Gaertnerhoerse und Gartenwelt*. 90:1528-1529.
- Grable, A.R. and E.G. Siemer. 1968. Effect of bulk density, aggregate size, and soil water suction on oxygen diffusion redox potentials, and elongation of corn roots. *Soil Sci. Soc. Am. Proc.* 32:180-186.
- Hendrix, P.F., C.R. Han, and P.M. Groffman. 1988. Soil respiration and no-tillage agroecosystems under different winter cover crop rotations. *Soil Till. Res.* 12:135-148.
- Hsu, J.T., D.B. Faulkner, K.A. Garleb, R.A. Barclay, G.C. Fahey, and L.L. Berger. 1987. Evaluation of corn fiber, cottonseed hulls, oat hulls and soybean hulls as roughage sources for ruminants. *J. Animal Sci.* 65(1):244-255..
- Huck, M.G. 1970. Variation in taproot elongation rate as influenced by composition of the soil air. *Agron. J.* 62:815-818
- Jayawardane, N.S. and W.S. Meyer. 1985. Measuring air-filled porosity changes in an irrigated swelling clay soil. *Aust. J. Soil Res.* 23:15-22.

- Jones, C. A. 1983. Effect of soil texture on critical bulk densities for root growth. *Soil Sci. Soc. Am. J.* 47:1208-1211.
- Logsdon, S.D. and C.A. Cambardella. 2000. Depth-incremental soil properties under no-till or chisel management. *Soil Sci. Soc. Am. J.* 63:197-200.
- Logsdon, S.D., T.C. Kaspar, and C.A. Cambardella. 1999. Temporal changes in small depth-incremental soil bulk density. *Soil Sci. Soc. Am. J.* 64:710-714.
- Marschner, Horst. 1986. *Mineral Nutrition of Higher Plants.* p.213. Academic Press Limited. San Diego, Calif.
- McAfee, M., J. Lindstrom, and W. Johansson. 1989. Aeration changes after irrigation in a clay soil. *J. Soil Sci.* 40:719-729.
- Meek, B.D., C.F. Ehlig, L.H. Stolzy, and L.E. Graham. 1983. Furrow and trickle irrigation: Effects on soil oxygen and ethylene and tomato yield. *Soil Sci. Soc. Am. J.* 47:631-635.
- Puentes, R., B.L. Harris, and C. Victora. 1988. Management of vertisols of temperate regions. pp.129-130. In: L.P. Wilding and R. Puentes (Eds). *Vertisols: Their Distribution, Properties, Classification and Management.* Tech. Monograph No. 18. Soil Management Support Services. Texas A&M Univ. Printing Center, College Station, Tex.
- Salisbury, F.B. and C.W. Ross. 1985. *Plant Physiology*, 3rd Ed. pp. 114-116. Wadsworth, Inc Belmont, Calif.
- SAS Institute, Inc. 2001. *The SAS System for Windows.* Release 8.02. SAS Institute, Inc. Cary, N.C.

Sloan, J.J, W. Mackay, and S. George. 2000. Growing mediums for porous pavement and rooftop gardens. pp. 321-327. In: W. Burghardt and C. Dornauf (eds). Proc. First Intl. Conf. Soils of Urban, Industrial, Traffic and Mining Areas. Universität-GH Essen, Essen, Germany

Steffen, K. 1989. A hanging basket plant with future—*Scaevola aemula*. Gaertnermeister 92:944-945.

United States Golf Association, Green Section Staff. 1993. Specifications for a method of putting green construction. U.S.G.A., Far Hills, N.J.

Voorhees, W.B., D.A. Farrell, and W.E. Larson. 1975. Soil strength and aeration effects on root elongation. Soil Sci. Soc. Am. Proc. 39:948-953.

Table 1. Effect of soil amendments on survival of pansy and scaevola plants in a clay soil during two growing seasons.

Soil amendment	Plant survival rate (% of total planted)			
	Pansy ^y		Scaevola ^y	
	Dec. 1996 – June 1997	Nov. 1997 – June 1998	June 1997 – Oct. 1997	June 1998 – Dec. 1998
Control (Ctrl)	96 a ^z	71 a	8 b	58 b
Small dia. expanded shale (SSh)	95 a	95 a	30 b	80 a
Large dia. expanded shale (LSH)	93 a	86 a	79 a	92 a
Sand	100 a	96 a	17 b	17 b
Sphagnum peat moss (SPM)	100 a	83 a	50 ab	75 a
Cottonseed hulls (CH)	100 a	100 a	67 a	67 a

^z Means within a column followed by the same letter are not statistically different (Duncan's multiple range test, $p \leq 0.05$).

^y Values represent survival of total pansies (n=24) and scaevola (n=12) planted.




Figure 1. Monthly rainfall totals (A) and average monthly maximum temperatures (B) for 1997 and 1998.




Figure 2. Effects of small diameter expanded shale (SSh), large diameter expanded shale (LSh), sand, sphagnum peat moss (SPM), and cottonseed hulls (CH) on bulk density of Austin silty clay (Ctrl) soil seven months after application ($1000 \text{ kg m}^{-3} = 62 \text{ lb ft}^{-3}$). Columns with the same letter are not statistically different at the 0.05 level of probability, $n=6$.




Figure 3. Effects of small diameter expanded shale (SSh), large diameter expanded shale (LSh), sand, sphagnum peat moss (SPM), and cottonseed hulls (CH) on pansy foliage quality (A), number of blossoms per plant (B), pansy dry matter yield (C), pansy N content (D), and root weight per plant (E) compared to a non-amended Austin silty clay soil (Ctrl) for two growing seasons (1 lb = 454 g; 1 g kg⁻¹ = 0.1 percent). Columns within a growing season with the same letter are not statistically different (Duncan's multiple range test, P<0.05, n=6).




Figure 4. Effects of small diameter expanded shale (SSh), large diameter expanded shale (LSh), sand, sphagnum peat moss (SPM), and cottonseed hulls (CH) on scaevola foliage quality (A) and blossom coverage (B) ratings in a clay soil (Ctrl) for two growing seasons. Columns within a growing season with the same letter are not statistically different (Duncan's multiple range test, $P \leq 0.05$, $n=6$).